

# On the number of generators for transeunt triangles

J. T. Butler\*, G. W. Dueck†, S. N. Yanushkevich‡, V. P. Shmerko§

November 30, 1999

\*Dept. of Electrical and Computer Eng., Naval Postgraduate School, Code EC/Bu Monterey, CA 93943-5121, USA, butler@cs.nps.navy.mil

†Faculty of Comp. Sci, Univ. of New Brunswick, Fredericton, NB, E3B 5A3 CANADA, gdueck@unb.ca

‡ §Dept. of CAD Systems, Faculty of CS and IS, Technical Univ., Zolnierska str. 49, 71210 Szczecin, POLAND shmerko@beta.ii.tuniv.szczecin.pl., januszki@beta.ii.tuniv.szczecin.pl. / University of Informatics & Radioelectronics, Minsk, BELARUS

## Abstract

A transeunt triangle of size  $n$  consists of  $(n+1) \times (n+1) \times (n+1)$  0's and 1's whose values are determined by the sum modulo 2 of two other local values. For a given  $n$ , two transeunt triangles of size  $n$  can be combined using the element-by-element modulo 2 sum to generate a third transeunt triangle. We show that, for large  $n$ , the  $\frac{1}{3}2^{n+1}$  transeunt triangles of size  $n$  can be generated from a set of only  $\frac{n}{3}$  generator transeunt triangles.

**Index Terms.** Symmetric functions, Reed-Muller expansion, transeunt triangle.

## 1 Introduction

A transeunt triangle of size  $n$  is completely specified by a binary  $n+1$ -tuple that forms the first row. The second row, a binary  $n$ -tuple, is formed by the pairwise modulo 2 sum of adjacent elements in the first row. Similarly, elements of all other rows are specified as the modulo 2 sum of adjacent elements in the row above it. The last row is a single 0 or 1. Fig. 1 shows all transeunt triangles of size 3. Here, a transeunt triangle is not repeated if it is the same as the rotation of another triangle in the figure.

---

\*Research supported by NPS Direct Grant Funded UHCD1 (USA)

†Research supported by the NSERC (CANADA)

‡Research supported in part from the St. Francis Xavier University (CANADA), in part by Committee of Scientific Researchers (POLAND) and in part by the Foundation of Fundamental Research, National Academy of Sciences (BELARUS)

§The author gratefully acknowledges the facilities provided as *James Chair Professor* from the St. Francis Xavier University (CANADA)

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>30 NOV 1999</b>		2. REPORT TYPE		3. DATES COVERED	
4. TITLE AND SUBTITLE <b>On the number of generators for transeunt triangles</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Naval Postgraduate School, Department of Electrical and Computer Engineering, Monterey, CA, 93943</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited.</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT <b>A transeunt triangle of size n consists of (n+1)?(n+1)?(n+1) 0's and 1's whose values are determined by the sum modulo 2 of two other local values. For a given n two transeunt triangles of size n can be combined using the element-by-element modulo 2 sum to generate a third transeunt triangle. We show that, for large n, the 1 32n+1 transeunt triangles of size n can be generated from a set of only n 3 generator transeunt triangles.</b>					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES <b>8</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			




Figure 1: All transeunt triangles of size 3.

Note that a transeunt triangle can be formed from *any* of its three sides. That is, if instead one begins with a side produced by the process described above, a rotated version of the same transeunt triangle results.

The transeunt triangle is the basis of an efficient algorithm for determining the minimal fixed polarity Reed-Muller canonical expansion of a totally symmetric switching function [1], [3], [4]. An  $n$ -variable totally symmetric function is specified by a binary  $n + 1$ -tuple that becomes one of the triangle's sides. Certain rectangles within the transeunt triangle each represent a circuit implementation of the symmetric function described by the edge. Transeunt triangles are related to the Sierpiński's Gasket [2]. If the first row has exactly one 1, a Sierpiński's Gasket forms below that 1. This embedded triangle is Pascal's triangle modulo 2. Transeunt triangles are also related to a specific one-dimensional cellular automata system. In such systems, the next state (0 or 1) of every cell is the modulo 2 sum of its present state and the present state of its right neighbor. This particular system is Rule 102 in Wolfram's classification scheme [5], which belongs to the class of *additive* cellular automata systems. The transeunt triangle shows the sequence of states in such a cellular automata system, where the first row is the initial state.

We are interested in determining the minimum number of generators of all transeunt triangles of size  $n$ . That is, given two transeunt triangles, a third can be generated by forming the bit-by-bit modulo 2 sum. For example, forming the bit-by-bit modulo 2 sum of any triangle with itself generates **0**, the all 0 transeunt triangle. All eight transeunt triangles

$$\begin{array}{cccc}
1 & 1 & 1 & 1 \\
0 & 0 & 0 & \\
0 & 0 & & \\
0 & & & 
\end{array}
\quad
\begin{array}{cccc}
0 & 1 & 0 & 0 \\
1 & 1 & 0 & \\
0 & 1 & & \\
1 & & & 
\end{array}$$

Figure 2: Generator transeunt triangles for all transeunt triangles of size 3.

of size 3 in Fig. 1 can be formed as some combination, including rotated versions, of the two transeunt triangles shown in Fig. 2. Since one transeunt triangle is not sufficient to produce all others, the minimum number of generators needed is 2. In the general case, all transeunt triangles can be generated from transeunt triangles with a single 1 on an edge. Therefore, it is sufficient to have  $n + 1$  generator transeunt triangles (or even less considering transeunt triangles where two different binary  $n + 1$ -tuples with exactly one 1 occur as sides). However, one can do better.

**Definition 1.1.** Let  $g_{\text{approx}}(n)$  and  $g_{\text{exact}}(n)$  be two functions of  $n$ .  $g_{\text{exact}}(n) \sim g_{\text{approx}}(n)$  if and only if

$$\lim_{n \rightarrow \infty} \frac{g_{\text{approx}}(n)}{g_{\text{exact}}(n)} = 1. \quad (1.1)$$

In what follows,  $g_{\text{approx}}(n)$  is a *simple*, approximate expression for  $g_{\text{exact}}(n)$ .

**Theorem 1.1.** The number  $N_{\text{tt}}(n)$  of unoriented transeunt triangles of size  $n$  is

$$N_{\text{tt}}(n) \sim \frac{1}{3} 2^{n+1}, \quad (1.2)$$

and the minimum number  $\tau_{\text{tt}}(n)$  of (generator) transeunt triangles needed to generate these triangles is

$$\tau_{\text{tt}}(n) \sim \frac{n}{3}. \quad (1.3)$$

## 2 Number of transeunt triangles and self-similar transeunt triangles

In this section, we prove the first part of Theorem 1.1, namely (1.2). Note that four of the transeunt triangles in Fig. 1 (those in the left column) are unchanged by a rotation of  $120^\circ$  and  $240^\circ$ .

**Definition 2.1.** A transeunt triangle  $T$  is *self-similar* if and only if a rotation of  $120^\circ$  and  $240^\circ$  leaves  $T$  unchanged.

**Theorem 2.1.** The number  $N_{\text{ss}}(n)$  of self-similar transeunt triangles of size  $n$  is

$$N_{\text{ss}}(n) = 2^{\lfloor \frac{n-1}{3} \rfloor + (n-1) \bmod 3}. \quad (2.1)$$

**Proof** For  $n = 1, 2$ , and  $3$ , the number of self-similar transeunt triangles is 1 ( $[00]$ ), 2 ( $[000]$  &  $[010]$ ), and 4 ( $[0000]$ ,  $[0110]$ ,  $[1101]$ , &  $[1011]$ ), respectively. For  $n = 1, 2$ , and  $3$ , (2.1) yields 1, 2, and 4, respectively. Consider a self-similar transeunt triangle of size  $n > 3$ . It embeds another triangle  $T'$  of size  $(n-2) \times (n-2) \times (n-2)$ , where  $n-2 > 1$ . Since  $T$  is invariant under rotation, so also is  $T'$ .

Given any  $(n-2) \times (n-2) \times (n-2)$  self-similar triangle  $T'$ , there are two ways to form an  $(n+1) \times (n+1) \times (n+1)$  self-similar triangle  $T$  (i.e. two circumferences that are complements). Thus,

$$N_{ss}(n) = 2N_{ss}(n-3). \quad (2.2)$$

But, this implies (2.1), since substituting the inductive hypothesis,

$$N_{ss}(n-3) = 2^{\lfloor \frac{n-4}{3} \rfloor + (n-4) \bmod 3}, \quad (2.3)$$

into (2.2), yields (2.1). ■

**Theorem 2.2.** *The number of unoriented transeunt triangles of size  $n$  is*

$$N_{tt}(n) = \frac{2^{n+1} + 2^{\lfloor \frac{n+2}{3} \rfloor + (n-1) \bmod 3}}{3}. \quad (2.4)$$

**Proof** Among the three sides in a transeunt triangle of size  $n$  are one, two, or three distinct binary  $n+1$ -tuples. As shown in Fig. 1, one and three are possible. For example, a transeunt triangle with one tuple is self-similar. Transeunt triangles with exactly two distinct tuples are impossible because of symmetry. It follows that the number of unoriented transeunt triangles is

$$N_{tt}(n) = \frac{2^{n+1} - N_{ss}(n)}{3} + N_{ss}(n) = \frac{2^{n+1} + 2N_{ss}(n)}{3}. \quad (2.5)$$

Substituting (2.3) into (2.5) yields (2.4) ■

Note that  $N_{tt}(n) \sim \frac{2^{n+1}}{3}$ . As the number  $n$  of variables increases without bound, the number of unoriented transeunt triangles approaches one-third the number of oriented transeunt triangles. This proves the first part of Theorem 1.1. This shows that most large transeunt triangles are *not* self-similar.

### 3 Generators of transeunt triangles

In this section, we prove the second part of Theorem 1.1, namely (1.3). Given a transeunt triangle  $T$ ,  $T_{120}$  and  $T_{240}$  are  $T$  rotated by  $120^\circ$  and  $240^\circ$ , respectively. Let  $\oplus$  be the bit-by-bit modulo 2 sum of two transeunt triangles.

**Lemma 3.1.** *Given any transeunt triangle  $T$ ,  $T \oplus T_{120} \oplus T_{240}$  is a self-similar transeunt triangle.*

**Definition 3.1.** Transeunt triangle  $T$  is a  $T_{ss}$ -self-similar-group transeunt triangle, where  $T_{ss} = T \oplus T_{120} \oplus T_{240}$ .

Since  $T_{ss} \oplus T_{ss120} \oplus T_{ss240} = T_{ss}$ ,  $T_{ss}$  is a  $T_{ss}$ -self-similar-group transeunt triangle. Further, there is exactly one self-similar  $T_{ss}$ -self-similar-group transeunt triangle, namely  $T_{ss}$ .

**Definition 3.2.**  $T$  is a *basic transeunt triangle* if it is a  $T_{ss}$ -self-similar-group transeunt triangle, where  $T_{ss} = \mathbf{0}$ , the transeunt triangle with all 0 entries.

Given transeunt triangle  $T$ ,  $T' = T \oplus T_{ss}$  is a basic transeunt triangle, since  $T' \oplus T'_{120} \oplus T'_{240} = (T \oplus T_{ss}) \oplus (T_{120} \oplus T_{120ss}) \oplus (T_{240} \oplus T_{240ss}) = T_{ss} \oplus T_{ss} = \mathbf{0}$ . Further,  $T' = T \oplus T \oplus T_{120} \oplus T_{240} = T_{120} \oplus T_{240} = T \oplus T_{120}$ .

**Definition 3.3.** Transeunt triangle  $T$  is a  $T_b$ -basic-group transeunt triangle, where  $T_b = T \oplus T_{120}$ .

**Theorem 3.2.** *The number of unoriented basic transeunt triangles of size  $n$  is*

$$N_b(n) = \frac{4^{\lceil \frac{n}{3} \rceil} + 2}{3}. \quad (3.1)$$

**Proof** For every unoriented basic transeunt triangle  $T'$ , there is a unique triangle in self-similar-group  $T_{ss}$ , that is obtained as  $T' \oplus T_{ss}$ . Similarly, for every unoriented transeunt triangle  $T''$  in self-similar-group  $T_{ss}$ , there is a unique unoriented basic transeunt triangle, that is obtained as  $T'' \oplus T_{ss}$ . Thus, each self-similar-group represents an equal sized block in a partition of all unoriented transeunt triangles. Therefore, the number of unoriented basic transeunt triangles is

$$N_b(n) = \frac{N_{tt}(n)}{N_{ss}(n)}. \quad (3.2)$$

Substituting (2.4) and (2.1) into (3.2) yields (3.1). ■

From Theorem 3.2,  $N_b(n) \sim \frac{1}{3}4^{\lceil \frac{n}{3} \rceil}$ . From Definitions 3.1 and 3.3, it follows that each transeunt triangle  $T$  is characterized by the self-similar transeunt triangle  $T_{ss}$  and by the basic transeunt triangle  $T_b$ , and that

$$T = T_{ss} \oplus T_b. \quad (3.3)$$

Therefore, if we can generate all self-similar and all basic transeunt triangles, we can generate all transeunt triangles. Both the set of basic transeunt triangles and the set of self-similar transeunt triangles are closed under the  $\oplus$  operation. Thus,  $T'' = T \oplus T'$  is a basic transeunt triangle if  $T$  and  $T'$  are basic, and  $T''$  is self-similar if  $T$  and  $T'$  are self-similar. Thus, each set can be generated from a subset. Indeed,

**Theorem 3.3.** *The set of self-similar transeunt triangles of size  $n > 1$  can be generated from a set of  $\alpha(n)$  nonredundant self-similar generator transeunt triangles, where*

$$\alpha(n) = \left\lfloor \frac{n-1}{3} \right\rfloor + (n-1) \bmod 3. \quad (3.4)$$

**Proof** There exists a set  $S_{ss} = \{T_1, T_2, \dots, T_m\}$  of self-similar transeunt triangles from which all self-similar transeunt triangles can be generated as the modulo 2 sum of elements from  $S_{ss}$  (e.g.  $S_{ss}$  can be all self-similar transeunt triangles). If  $S_{ss}$  is nonredundant, then no  $T_i \in S$  is the modulo 2 sum of any other transeunt triangles in  $S$ . Since  $T_i \oplus T_i = \mathbf{0}$ , no element in  $S$  is chosen more than once. It follows that  $m \geq \left\lfloor \frac{n-1}{3} \right\rfloor + (n-1) \bmod 3$ ; otherwise, there are not enough combinations ( $2^m$ ) of transeunt triangles in  $S$  to form all  $2^{\left\lfloor \frac{n-1}{3} \right\rfloor + (n-1) \bmod 3}$  self-similar transeunt triangles (for convenience, we substitute  $\mathbf{0}$  for the combination consisting of choosing *no* element of  $S$ ). We show that equality holds by showing that the modulo 2 sum of a combination of elements from  $S$  is distinct from any other combination. On the contrary, suppose

$$T_{i_1} \oplus T_{i_2} \oplus \dots \oplus T_{i_\alpha} \oplus T_{j_1} \oplus T_{j_2} \oplus \dots \oplus T_{j_\beta} = T_{k_1} \oplus T_{k_2} \oplus \dots \oplus T_{k_\gamma} \oplus T_{j_1} \oplus T_{j_2} \oplus \dots \oplus T_{j_\beta}, \quad (3.5)$$

where the only  $T$ 's common to both sides of (3.5) are  $T_{j_1}, T_{j_2}, \dots$ , and  $T_{j_\beta}$ . It follows that

$$T_{i_1} \oplus T_{i_2} \oplus \dots \oplus T_{i_\alpha} = T_{k_1} \oplus T_{k_2} \oplus \dots \oplus T_{k_\gamma}, \quad (3.6)$$

and that

$$T_{i_1} = T_{k_1} \oplus T_{k_2} \oplus \dots \oplus T_{k_\gamma} \oplus T_{i_2} \oplus \dots \oplus T_{i_\alpha}. \quad (3.7)$$

But, this is impossible, as no element of  $S$  is the modulo 2 sum of other elements of  $S$ . The theorem statement follows. ■

**Theorem 3.4.** *The set of unoriented basic transeunt triangles of size  $n$  can be generated from a set of  $\beta(n)$  nonredundant unoriented basic generator transeunt triangles, where*

$$\beta(n) = \left\lceil \frac{n}{3} \right\rceil. \quad (3.8)$$

**Proof** There exists a set  $S_b = \{T_1, T_2, \dots, T_p\}$  of basic generator transeunt triangles from which all basic transeunt triangles can be generated as the modulo 2 sum of elements from  $S_b$ . The all 0 triangle can be generated as the element-by-element modulo 2 sum of any non-zero generator triangle with itself. The remaining  $(4^{\left\lceil \frac{n}{3} \right\rceil} - 1)/3$  non-zero basic transeunt triangles can be generated as follows. Each generator transeunt triangle  $T$  can be used in four ways 1. omitted, 2. as  $T$ , 3. as  $T_{120}$ , and 4. as  $T_{240}$ . For each combination of at least

one generator, rotating *all* generator triangles by  $120^\circ$  or by  $240^\circ$  creates a resulting triangle that is rotated by  $120^\circ$  or  $240^\circ$ , respectively; however this produces the *same* unoriented triangle. This accounts for the divisor of 3 in  $(4^{\lceil \frac{n}{3} \rceil} - 1)/3$ . The one exception, where all generator triangles are omitted, accounts for the  $-1$  in  $(4^{\lceil \frac{n}{3} \rceil} - 1)/3$ . The argument regarding uniqueness of combinations is similar to that of Theorem 3.3. ■

For  $S_b$  and  $S_{ss}$ , the set of basic and self-similar generator transeunt triangles, respectively, we can form a set  $S_{b/ss} = \{T_{b/ss} = T_b \oplus T_{ss} | T_b \in S_b, T_{ss} \in S_{ss}\}$ . We can then form every basic generator transeunt triangle and every self-similar transeunt triangle, as  $T_{b/ss120} \oplus T_{b/ss240}$  and  $T_{b/ss} \oplus T_{b/ss120} \oplus T_{b/ss240}$ , respectively. From the generator triangles, we can form all basic and all self-similar transeunt triangles. This proves the following.

**Theorem 3.5.** *All basic and all self-similar transeunt triangles can be generated from  $\gamma(n)$  generator transeunt triangles, where*

$$\gamma(n) = \max \left\{ \left\lceil \frac{n}{3} \right\rceil, \left\lfloor \frac{n-1}{3} \right\rfloor + (n-1) \bmod 3 \right\}. \quad (3.9)$$

From all basic and all self-similar transeunt triangles, we can generate all transeunt triangles. Note that  $\gamma(n)$  is also the *minimum* number of transeunt triangles from which all transeunt triangles can be generated. On the contrary, with fewer than  $\gamma(n)$  generators, it is impossible to generate either all basic and/or all self-similar transeunt triangles, as observed in the proofs of Theorems 3.3 and 3.4. Observing that  $\gamma(n)$  approaches  $\frac{n}{3}$  as  $n$  increases without bound proves the second part of Theorem 1.1. Table 1 shows the number of transeunt triangle, the number of self-similar transeunt triangles, and the number of basic transeunt triangles, for  $1 \leq n \leq 20$ .

## 4 Concluding remarks

In this paper, we have shown that the minimum number of generators of transeunt triangles is small, approaching  $\frac{n}{3}$  as the size  $n$  increases without bound. An appreciation for the reduction achievable can be obtained by an examination of Table 1, which shows, for example, that the number of transeunt triangles of size  $n = 20$  is 699,136, while the minimum number of generators for this set is only 7.

## References

- [1] J. T. Butler, G. W. Dueck, V. P. Shmerko, and S. N. Yanushkevich. An efficient algorithm for the synthesis of fixed polarity Reed-Muller expansions of symmetric functions. *preprint*, 1999.



$n$	Number of Transeunt Triangles			Number of Generators	
	Total $N_{tt}(n)$	Self-Similar $N_{ss}(n)$	Unoriented Basic $N_b(n)$	Self-Similar $\alpha(n)$	Unoriented Basic $\beta(n)$
1	2	1	2	0	1
2	4	2	2	1	1
3	8	4	2	2	1
4	12	2	6	1	2
5	24	4	6	2	2
6	48	8	6	3	2
7	88	4	22	2	3
8	176	8	22	3	3
9	352	16	22	4	3
10	688	8	86	3	4
11	1,376	16	86	4	4
12	2,752	32	86	5	4
13	5,472	16	342	4	5
14	10,944	32	342	5	5
15	21,888	64	342	6	5
16	43,712	32	1,366	5	6
17	87,424	64	1,366	6	6
18	174,848	128	1,366	7	6
19	349,568	64	5,462	6	7
20	699,136	128	5,462	7	7
$\infty$	$\frac{1}{3}2^{n+1}$	$2^{\lfloor \frac{n-1}{3} \rfloor}$	$\frac{1}{3}4^{\lceil \frac{n}{3} \rceil}$	$\frac{n}{3}$	$\frac{n}{3}$

Table 1: Number of  $n$ -variable transeunt triangles, self-similar, basic unoriented, generator self-similar, and generator basic transeunt triangles, for  $1 \leq n \leq 20$ .

- [2] I. Stewart. Four encounters with Sierpiński's Gasket. *J. Mathematical Intelligencer*, 17(1):52–64, 1999.
- [3] V. P. Suprun. Polynomial expression of symmetric Boolean functions. *Izvestija AN USSR. Techn. Kibernetika (in Russian)*, (4):123–127, 1985.
- [4] V. P. Suprun. Fixed polarity Reed-Muller expressions of symmetric Boolean functions. In *Proc. IFIP WG 10.5 Workshop on Application of the Reed-Muller Expansions in Circuit Design*, pages 246–249, 1995.
- [5] S. Wolfram. Theory and applications of cellular automata. *World Scientific Publishing Co. Ltd.*, pages 485–557, 1986.